

Chadwick B. Hampton, MD

CONSENT FOR JOINT INJECTION

As with any injection, you will feel pain. The use of an ice pack applied to the injection site two to three times a day for the next two to four days should help minimize any discomfort you may feel.

The use of an over the counter medication, such as *Tylenol*, *Advil* or *Aleve* can be taken to lessen the discomfort. Please read the back to these medication bottles for directions. They can be taken up to one week after the injection. In some cases, your doctor may prescribe a medicine that you may also take instead of the above-mentioned medications.

The severity of your condition can determine how much relief you will receive from the injection. There is no guarantee that the injection will help your condition. You may feel relief for hours, days, weeks or even months. Every patient is different. You should also avoid activities that put stress on the joint for the first four days after the injection.

3400 Burns Road Suite 101

Palm Beach Gardens, FL 33410

Tel: 561.327.9116

Tel: 561.588.9912

TEAMHAMPTON@FLORTHOCARE.COM



1049 South State Road 7

Wellington, FL 33414

Tel: 561.327.9116

Tel: 561.588.9912

TEAMHAMPTON@FLORTHOCARE.COM

Thank you,

Dr. Hampton

Patient Name: _____ **DOB:** _____

Patient Signature: _____ **Witness:** _____

Account No: _____ **Date:** _____